

®

Dried Smallpox Vaccine

only

**DO NOT INJECT INTRAMUSCULARLY (IM), INTRAVENOUSLY (IV), OR
SUBCUTANEOUSLY (SC). FOR CONVENTIONAL SMALLPOX VACCINATION
(SCARIFICATION) ONLY.**

DESCRIPTION

reaction is passed within three days following the application of fully potent vaccine with an antibody rise occurring in roughly half of those who exhibit such a reaction. As immunity wanes, revaccination with potent vaccine elicits this allergic response followed by the changes produced by propagating virus. The lesion may then go through the same course as the primary vaccination

PRECAUTIONS
General

The most important measure to prevent inadvertent implantation and contact transmission from vaccinia vaccination is thorough hand washing after changing the bandage or after any other contact with the vaccination site.

ADVERSE REACTIONS

The United States Department of Health and Human Services (DHHS) has established the Vaccine Adverse Event Reporting System (VAERS) to accept all reports of suspected adverse

6. Remove rubber cover from end of the diluent syringe.

13. Record date of reconstitution.

5. Peel back the packaging approximately halfway exposing the butt-end of needle.



14. Repeat Steps 3 through 10.

15. If vaccine is to be restored for subsequent use, replace stopper-cap and store at 2° to 8°

HOW SUPPLIED

Combination package of 1 vial of Dried Smallpox Vaccine, 1 Diluent syringe (0.25 mL), 1 vented needle, 100 individually wrapped bifurcated needles (20 strips, 5 needles per strip).

10. Vaccinia outbreak – Nevada.